

**KURZPROTOKOLL**  
**MK7902-005**

<b>Öffentlicher Titel</b>	Phase II Studie zu Lenvatinib und Pembrolizumab bei vorbehandelten soliden Tumoren
<b>Wissenschaftl. Titel</b>	A Multicenter, Open-label Phase 2 Study of Lenvatinib (E7080) Plus Pembrolizumab (MK-3475) in Previously Treated Subjects with Selected Solid Tumors
<b>Kurztitel</b>	MK7902-005
<b>Studienart</b>	multizentrisch, prospektiv, Therapiestudie, offen/unverblindet, einarmig, Pharma-Studie
<b>Studienphase</b>	Phase II
<b>Erkrankung</b>	Nervensystem: Gliome: Glioblastom (WHO Grad IV) - Zweitlinie oder höher
<b>Einschlusskriterien</b>	<ul style="list-style-type: none"><li>- Have a histologically or cytologically-documented, advanced (metastatic and/or unresectable) solid tumor that is incurable and for which prior standard systemic therapy has failed in one of the following cohorts: Cohort E: Glioblastoma</li><li>- Participants must have progressed on or since the last treatment</li><li>- Have measurable disease per RANO for the GBM</li><li>- Life expectancy of 12 weeks or more</li><li>- (ECOG) performance status of 0 to 1 within 7 days of treatment initiation</li><li>- Adequately controlled blood pressure (BP) with or without antihypertensive medications, defined as BP <math>\leq</math>150/90 mm Hg</li></ul>
<b>Ausschlusskriterien</b>	<ul style="list-style-type: none"><li>- Prolongation of QTc interval (calculated using Fridericia's formula) to <math>&gt;</math>480 ms.</li><li>- Has left ventricular ejection fraction (LVEF) <math>&lt;</math>55 as determined by multigated acquisition scan (MUGA) or echocardiogram (ECHO).</li><li>- Has a diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy (in dosing exceeding 10 mg daily of prednisone equivalent) or any other form of immunosuppressive therapy within 7 days prior the first dose of study drug.</li><li>- HIV, Hep B, Hep C, Tuberculosis</li><li>- Has carcinomatous meningitis</li><li>- Has recurrent tumor greater than 6cm in maximum diameter.</li><li>- The GBM participant will not be excluded from the study for systemic steroid therapy, as long as dexamethasone or its steroid equipotent dosing equivalent is administered at a constant dose for at least 2 weeks prior to study treatment start where the dexamethasone dose (or its equivalent) is <math>\leq</math> 2 mg daily.</li></ul>
<b>Alter</b>	18 Jahre und älter
<b>Prüfzentren</b>	<b>Neuroonkologie</b> (Nachbeobachtung) Schleusenweg 2-16 60528 Frankfurt am Main Dr. med. Michael Burger Tel: 069 6301-87711 Fax: 069 6301-87713 <a href="mailto:michael.burger@unimedizin-ffm.de">michael.burger@unimedizin-ffm.de</a> <b>Universitätsklinikum Frankfurt</b> (Nachbeobachtung) Dr. Senckenbergisches Institut für Neuroonkologie Theodor-Stern-Kai 7 60590 Frankfurt am Main Dr. med. Michael Burger Tel: 069 6301-87711 Fax: 069 6301-87713 <a href="mailto:michael.burger@unimedizin-ffm.de">michael.burger@unimedizin-ffm.de</a>
<b>Sponsor</b>	MSD Sharp & Dohme
<b>Registrierung in anderen Studienregistern</b>	ClinicalTrials.gov NCT03797326 (primäres Register) EudraCT 2018-003747-37