

## **KURZPROTOKOLL** **AMLSG 15-10**

<b>Öffentlicher Titel</b>	Cytarabin und Etoposid mit oder ohne ATRA bei AML mit NPM1-Mutation
<b>Wissenschaftl. Titel</b>	Randomized Phase III Study of Low-Dose Cytarabine and Etoposide with or without All-Trans Retinoic Acid in Older Patients not Eligible for Intensive Chemotherapy with Acute Myeloid Leukemia and NPM1 Mutation
<b>Kurztitel</b>	AMLSG 15-10
<b>Studienart</b>	multizentrisch, Therapiestudie, randomisiert, offen/unverblindet, zweiarmig, Investigator Initiated Trial (IIT)
<b>Studienphase</b>	Phase III
<b>Erkrankung</b>	Blut: Akute myeloische Leukämie (AML): Neu diagnostiziert / de novo
<b>Einschlusskriterien</b>	<ul style="list-style-type: none"><li>- Patients with confirmed diagnosis of acute myeloid leukemia according to the World Health Organization (WHO) classification (including de novo AML, t-AML and s-AML)</li><li>- Presence of NPM1 mutation as assessed in one of the central AMLSG reference laboratories.</li><li>- Age &gt; 60 years. There is no upper age limit.</li><li>- No prior chemotherapy for leukemia except hydroxyurea to control hyperleukocytosis if needed for up to 10 days during the diagnostic screening phase.</li><li>- Signed written informed consent</li><li>- Men must give their informed consent that they do not father a baby and must use a latex condom during any sexual contact with women of childbearing potential, even if they have undergone a successful vasectomy while on therapy and for 3 month after the last dose of chemotherapy.</li><li>- WHO performance status &lt;= 3</li><li>- Patients not eligible for intensive chemotherapy according to at least one of the following criteria</li><li>- HCT-CI Score &gt;2 (see Appendix F)</li><li>- Patient's decision</li></ul>
<b>Ausschlusskriterien</b>	<ul style="list-style-type: none"><li>- All other AML subtypes, in particular those AML with other recurrent genetic changes (according to WHO 2008):</li><li>- AML with t(8;21)(q22;q22); RUNX1-RUNX1T1</li><li>- AML with inv(16)(p13.1q22) or t(16;16)(p13.1;q22); CBFβ-MYH11</li><li>- AML with t(15;17)(q22;q12); PML-RARA (or other translocations involving RARA)</li><li>- AML with t(9;11)(p22;q23); MLLT3-MLL (or other translocations involving MLL)</li><li>- AML with t(6;9)(p23;q34); DEK-NUP214</li><li>- AML with inv(3)(q21q26.2) or t(3;3)(q21;q26.2); RPN1-EV11</li><li>- No consent for registration, storage and processing of the individual disease-characteristics and course as well as information of the family physician and all other treating physicians about study participation</li><li>- Bleeding disorder independent of leukemia</li><li>- Uncontrolled infection</li><li>- Known positive for HIV, HBV or HCV</li><li>- Organ insufficiency (creatinine &gt;1.5x upper normal serum level; bilirubin, AST or ALP &gt;2.5x upper normal)</li><li>- Severe neurological or psychiatric disorder interfering with ability of giving an informed consent</li><li>- Patients with a "currently active" second malignancy other than non-melanoma skin cancers. Patients are not considered to have a "currently active" malignancy if they have completed therapy and are considered by their physician to be at less than 30% risk of relapse within one year.</li></ul>

**KURZPROTOKOLL**  
**AMLSG 15-10**

<b>Alter</b>	> 60 Jahre
<b>Molekularer Marker</b>	NPM1
<b>Prüfzentren</b>	<b>MVZ-Osthessen GmbH</b> (Rekrutierung beendet) Onkologisches Zentrum Pacelliallee 4 36043 Fulda Dr. med. Andrea Distelrath Tel: 0661 845487 Fax: 0661 845484 <a href="mailto:roswitha.rausch@klinikum-fulda.de">roswitha.rausch@klinikum-fulda.de</a>
<b>Sponsor</b>	Universität Ulm
<b>Registrierung in anderen Studienregistern</b>	ClinicalTrials.gov NCT01237808 EudraCT 2010-023409-37