

KURZPROTOKOLL **GMALL-MOLACT1-BLINA**

Öffentlicher Titel	Blinatumomab bei MRD-positiver B-Vorläufer ALL
Wissenschaftl. Titel	A multicenter, single-arm study to assess the efficacy, safety, and tolerability of the BiTE® antibody blinatumomab in adult patients with minimal residual disease (MRD) of B-precursor acute lymphoblastic leukemia (Blast Successor Trial)
Kurztitel	GMALL-MOLACT1-BLINA
Studienart	multizentrisch, prospektiv, Therapiestudie, offen/unverblindet, einarmig, Investigator Initiated Trial (IIT)
Studienphase	Phase II
Erkrankung	Blut: Akute lymphatische Leukämie (ALL): Rezidiert/refraktär
Einschlusskriterien	<ul style="list-style-type: none">- Patients with CD19 positive B-precursor ALL in complete hematological remission defined as less than 5% blasts in bone marrow after at least three intense chemotherapy blocks (e.g., GMALL induction I-II/consolidation I).- Presence of minimal residual disease (MRD) at a level of $\geq 10^{-4}$ (molecular failure or molecular relapse) in an assay with a minimum sensitivity of 10^{-4} documented after an interval of at least 2 weeks from last systemic chemotherapy- For evaluation of MRD patients must have at least one molecular marker based on individual rearrangements of immunoglobulin, TCR-genes or other suitable genes evaluated by the reference laboratory of the trial- Bone marrow function as defined below: (a) ANC (Neutrophils) $\geq 1,000/\mu\text{L}$; (b) Platelets $\geq 50,000/\mu\text{L}$ (transfusion permitted); (c) HB level $\geq 9\text{g/dl}$ (transfusion permitted)- Renal and hepatic function as defined below: (a) AST (GOT), ALT (GPT), and AP < 5 x upper limit of normal (ULN); (b) Total bilirubin < 1.5 x ULN (unless related to Gilbert's Meulengracht disease); (c) Creatinine < 1.5 x ULN; (d) Creatinine clearance ≥ 60 mL/min (e.g. calculated according Cockcroft&Gault)- Negative HIV test, negative hepatitis B (HbsAg) and hepatitis C virus (anti-HCV) test- Negative pregnancy test in women of childbearing potential- ECOG Performance Status 0 or 1- Age ≥ 18 years- Ability to understand and willingness to sign a written informed consent- Signed and dated written informed consent is available- Participation in the registry of the German Multicenter Study Group for Adult ALL (GMALL)
Ausschlusskriterien	<ul style="list-style-type: none">- Ph/BCR-ABL positive ALL- Presence of circulating blasts or current extramedullary involvement by ALL- History or presence of clinically relevant CNS pathology (e.g. seizure, paresis, aphasia, cerebrovascular ischemia/hemorrhage, severe brain injuries, dementia, Parkinson's disease, cerebellar disease, organic brain syndrome or psychosis)- Current detection of ALL blast cells in cerebro-spinal fluid- History of or active relevant autoimmune disease- Systemic cancer chemotherapy within 2 weeks prior to study treatment (except for intrathecal prophylaxis)- Radiotherapy within 4 weeks prior to study treatment- Live vaccination within 2 weeks before the start of study treatment- Autologous hematopoietic stem cell transplantation (SCT) within six weeks prior to study treatment- Allogeneic SCT within 12 weeks before the start of study treatment

KURZPROTOKOLL GMALL-MOLACT1-BLINA

- Any active acute Graft-versus-Host Disease (GvHD), grade 2-4 according to the Glucksberg criteria or active chronic GvHD requiring systemic treatment
- Any systemic therapy against GvHD within 2 weeks before start of study treatment
- Therapy with monoclonal antibodies (rituximab, alemtuzumab) within 4 weeks prior to study treatment
- Treatment with any investigational product within four weeks prior to study treatment
- Previous treatment with blinatumomab or other anti-CD19-therapy
- Known hypersensitivity to immunoglobulins or to any other component of the study drug formulation
- History of malignancy other than ALL diagnosed within 5 years prior to start of protocol-specified therapy with the exception of: (a) Adequately treated non-melanoma skin cancer or lentigo maligna without evidence of disease; (b) Adequately treated cervical carcinoma in situ without evidence of disease; (c) Adequately treated breast ductal carcinoma in situ without evidence of disease; (d) Prostatic intraepithelial neoplasia without evidence of prostate cancer
- Active infection, any other concurrent disease or medical condition that are deemed to interfere with the conduct of the study as judged by the investigator
- Nursing women
- Woman of childbearing potential and is not willing to use 2 highly effective methods of contraception while receiving study treatment and for an additional 3 months after the last dose of study treatment.
- Male who has a female partner of childbearing potential, and is not willing to use 2 highly effective forms of contraception while receiving study treatment and for at least an additional 3 months after the last dose of study treatment

Alter	18 Jahre und älter
Molekularer Marker	CD19
Fallzahl	30
Prüfzentren	Universitätsklinikum Gießen und Marburg, Standort Marburg (Rekrutierung beendet) Hämatologie, Onkologie und Immunologie Baldingerstraße 35043 Marburg Iris Lenser Tel: 06421 58 63429 Fax: 06421 58 62703 studien-onkologie@uni-marburg.de Universitätsklinikum Frankfurt (Rekrutierung beendet) Medizinische Klinik II, Hämatologie/Onkologie Theodor-Stern-Kai 7 60590 Frankfurt am Main Studienkoordination GMALL-Molact-1-Blina molact1-blina@med.uni-frankfurt.de
Sponsor	Goethe-Universität Frankfurt
Förderer	AMGEN GmbH
Registrierung in anderen Studienregistern	ClinicalTrials.gov NCT03109093 EudraCT 2015-000733-76
Links	Studiendokumente zum Download (roXtra) Zu den Ein- und Ausschlusskriterien